

学位論文審査結果の要旨

所 属	三重大学大学院医学系研究科 甲 生命医科学専攻 臨床医学系講座 成育医学分野	氏 名	プリンス バッフアー トント Prince Baffour Tonto
審 査 委 員	主 査 野阪哲哉 副 査 堀浩樹 副 査 竹内万彦		
<p>(学位論文審査結果の要旨)</p> <p>High prevalence of IgE sensitization to inactivated influenza vaccines, yet robust IgG4 responses, in a healthy pediatric population</p> <p>【主論文審査結果の要旨】</p> <p>著者らは論文において下記の内容を述べている。</p> <p>Background: Anaphylaxis following influenza vaccination is a rare but serious problem. The underlying immune responses are not well understood. This study elucidated the IgE and IgG antibody responses in healthy children and adolescents following inactivated influenza vaccines (IIVs).</p> <p>Methods: The efficacy and safety of quadrivalent IIV (QIV) and trivalent IIV (TIV) were compared in healthy subjects aged 0-18 years. Serum IIV-specific IgE, IgG and IgG4 levels (sIgE, sIgG, sIgG4) were measured with ImmunoCAP. Hemagglutinin inhibition (HI) assay was performed for each influenza virus subtype. Sera from earlier patients who developed anaphylaxis to different IIVs were similarly tested.</p> <p>Results: A total of 393 subjects were enrolled: 96 were 6 months -2 years old, 100 were 3-5 years old, 100 were 6-12 years old, and 97 were 13-18 years old. No anaphylaxis was observed. Generally, QIV and TIV induced similar antibody responses. IIV-sIgE levels rose significantly after vaccination in the 6m-2y and 3-5y groups, did not change in the 6-12y group, and decreased in the 13-18y</p>			

group. In contrast, the IIV-sIgG4/sIgE ratio increased significantly after vaccination in all age groups. Sensitized subjects had significantly higher HI titers and IIV-sIgG levels in the youngest age group and higher IIV-sIgG4 levels in all age groups compared with the non-sensitized. The IIV-sIgG4/sIgE ratio in 5 patients with anaphylaxis was significantly lower than in age-matched healthy subjects.

Conclusion: IIVs induce IgE sensitization in healthy children, but also robust IgG4 responses that may protect them from anaphylaxis.

小児へのインフルエンザワクチン接種後のIgE、IgG抗体反応を明らかにした。正常小児でもIgEは産生されるが、阻止抗体IgG4がそれを上回って産生された。一方、アナフィラキシー患者ではIgG4産生が低いことを示した。アナフィラキシー発症のメカニズムを示すデータとして、学術上極めて有益であり、この論文を学位論文として価値あるものと認めた。

Influenza and Other Respiratory Viruses 2022; 1-10

Published: September 9, 2022

doi: 10.1111/irv.13053

Prince Baffour Tonto | Mizuho Nagao | Shigeru Suga | Kiyosu Taniguchi |
Masahiro Hirayama | Tetsuo Nakayama | Takuji Kumagai | Takao Fujisawa |
the Influenza Vaccine Study Group